PROBLEM 6.11

KNOWN: Radial distribution of local convection coefficient for flow normal to a circular disk.

FIND: Expression for average Nusselt number.

SCHEMATIC:



ASSUMPTIONS: Constant properties. **ANALYSIS:** The average convection coefficient is

$$\overline{h} = \frac{1}{A_s} \int_{A_s} h dA_s$$

$$\overline{h} = \frac{1}{\pi r_o^2} \int_0^{r_o} \frac{k}{D} N u_o \left[1 + a \left(r/r_o \right)^n \right] 2\pi r dr$$

$$\overline{h} = \frac{k N u_o}{r_o^3} \left[\frac{r^2}{2} + \frac{a r^{n+2}}{(n+2) r_o^n} \right]_0^{r_o}$$

where Nu_0 is the Nusselt number at the stagnation point (r = 0). Hence,

$$\overline{\mathrm{Nu}}_{\mathrm{D}} = \frac{\overline{\mathrm{hD}}}{\mathrm{k}} = 2\mathrm{Nu}_{\mathrm{O}} \left[\frac{\left(\mathrm{r/r_{\mathrm{O}}} \right)^2}{2} + \frac{\mathrm{a}}{\left(\mathrm{n+2} \right)} \left(\frac{\mathrm{r}}{\mathrm{r_{\mathrm{O}}}} \right)^{\mathrm{n+2}} \right]_{\mathrm{O}}^{\mathrm{r_{\mathrm{O}}}}$$

$$\overline{\mathrm{Nu}}_{\mathrm{D}} = \mathrm{Nu}_{\mathrm{O}} \left[1 + 2\mathrm{a}/\left(\mathrm{n+2} \right) \right]$$

$$\overline{\mathrm{Nu}}_{\mathrm{D}} = \left[1 + 2\mathrm{a}/\left(\mathrm{n+2} \right) \right] 0.814 \mathrm{Re}_{\mathrm{D}}^{1/2} \mathrm{Pr}^{0.36}.$$

<

COMMENTS: The increase in h(r) with r may be explained in terms of the sharp turn which the boundary layer flow must make around the edge of the disk. The boundary layer accelerates and its thickness decreases as it makes the turn, causing the local convection coefficient to increase.

Excerpts from this work may be reproduced by instructors for distribution on a not-for-profit basis for testing or instructional purposes only to students enrolled in courses for which the textbook has been adopted. Any other reproduction or translation of this work beyond that permitted by Sections 107 or 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful.